

# Multivariate Normal Distribution

---

Max Turgeon

STAT 4690—Applied Multivariate Analysis

## Building the multivariate density i

- Let  $Z \sim N(0, 1)$  be a standard (univariate) normal random variable. Recall that its density is given by

$$\phi(z) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}z^2\right).$$

- Now if we take  $Z_1, \dots, Z_p \sim N(0, 1)$  independently distributed, their joint density is

## Building the multivariate density ii

$$\begin{aligned}\phi(z_1, \dots, z_p) &= \prod_{i=1}^p \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}z_i^2\right) \\ &= \frac{1}{(\sqrt{2\pi})^p} \exp\left(-\frac{1}{2} \sum_{i=1}^p z_i^2\right) \\ &= \frac{1}{(\sqrt{2\pi})^p} \exp\left(-\frac{1}{2} \mathbf{z}^T \mathbf{z}\right),\end{aligned}$$

where  $\mathbf{z} = (z_1, \dots, z_p)$ .

- More generally, let  $\mu \in \mathbb{R}^p$  and let  $\Sigma$  be a  $p \times p$  positive definite matrix.

## Building the multivariate density iii

- Let  $\Sigma = LL^T$  be the Cholesky decomposition for  $\Sigma$ .
- Let  $\mathbf{Z} = (Z_1, \dots, Z_p)$  be a standard (multivariate) normal random vector, and define  $\mathbf{Y} = L\mathbf{Z} + \mu$ . We know from last lecture that
  - $E(\mathbf{Y}) = LE(\mathbf{Z}) + \mu = \mu$ ;
  - $\text{Cov}(\mathbf{Y}) = L\text{Cov}(\mathbf{Z})L^T = \Sigma$ .
- To get the density, we need to compute the inverse transformation:

$$\mathbf{Z} = L^{-1}(\mathbf{Y} - \mu).$$

## Building the multivariate density iv

- The Jacobian matrix  $J$  for this transformation is simply  $L^{-1}$ , and therefore

$$\begin{aligned} |\det(J)| &= |\det(L^{-1})| \\ &= \det(L)^{-1} \quad (L \text{ is p.d.}) \\ &= \sqrt{\det(\Sigma)}^{-1} \\ &= \det(\Sigma)^{-1/2}. \end{aligned}$$

## Building the multivariate density $\nu$

- Plugging this into the formula for the density of a transformation, we get

$$\begin{aligned} f(y_1, \dots, y_p) &= \frac{1}{\det(\Sigma)^{1/2}} \phi(L^{-1}(\mathbf{y} - \mu)) \\ &= \frac{1}{\det(\Sigma)^{1/2}} \left( \frac{1}{(\sqrt{2\pi})^p} \exp\left(-\frac{1}{2}(L^{-1}(\mathbf{y} - \mu))^T L^{-1}(\mathbf{y} - \mu)\right) \right) \\ &= \frac{1}{\det(\Sigma)^{1/2} (\sqrt{2\pi})^p} \exp\left(-\frac{1}{2}(\mathbf{y} - \mu)^T (LL^T)^{-1}(\mathbf{y} - \mu)\right) \\ &= \frac{1}{\sqrt{(2\pi)^p |\Sigma|}} \exp\left(-\frac{1}{2}(\mathbf{y} - \mu)^T \Sigma^{-1}(\mathbf{y} - \mu)\right). \end{aligned}$$

## Example i

```
set.seed(123)

n <- 1000; p <- 2
Z <- matrix(rnorm(n*p), ncol = p)

mu <- c(1, 2)
Sigma <- matrix(c(1, 0.5, 0.5, 1), ncol = 2)
L <- t(chol(Sigma))
```

## Example ii

```
Y <- L %*% t(Z) + mu
```

```
Y <- t(Y)
```

```
colMeans(Y)
```

```
## [1] 1.016128 2.044840
```

```
cov(Y)
```

```
##           [,1]      [,2]
```

```
## [1,] 0.9834589 0.5667194
```


```
## [2,] 0.5667194 1.0854361
```



## Example iii

```
library(tidyverse)
Y %>%
  data.frame() %>%
  ggplot(aes(X1, X2)) +
  geom_density_2d()
```

## Example iv



## Example v

```
library(mvtnorm)
```

```
Y <- rmvnorm(n, mean = mu, sigma = Sigma)
```

```
colMeans(Y)
```

```
## [1] 0.9812102 1.9829380
```


```
cov(Y)
```

## Example vi

```
##           [,1]      [,2]  
## [1,] 0.9982835 0.4906990  
## [2,] 0.4906990 0.9489171
```

```
Y %>%  
  data.frame() %>%  
  ggplot(aes(X1, X2)) +  
  geom_density_2d()
```

## Example vii



## Other characterizations

There are at least two other ways to define the multivariate random distribution:

1. A  $p$ -dimensional random vector  $\mathbf{Y}$  is said to have a multivariate normal distribution if and only if every linear combination of  $\mathbf{Y}$  has a *univariate* normal distribution.
2. A  $p$ -dimensional random vector  $\mathbf{Y}$  is said to have a multivariate normal distribution if and only if its distribution maximises entropy over the class of random vectors with fixed mean  $\mu$  and fixed covariance matrix  $\Sigma$  and support over  $\mathbb{R}^p$ .

## Useful properties i

- If  $\mathbf{Y} \sim N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ ,  $A$  is a  $q \times p$  matrix, and  $b \in \mathbb{R}^q$ , then

$$A\mathbf{Y} + b \sim N_q(A\boldsymbol{\mu} + b, A\boldsymbol{\Sigma}A^T).$$

- If  $\mathbf{Y} \sim N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  then all subsets of  $\mathbf{Y}$  are normally distributed; that is, write
  - $\mathbf{Y} = \begin{pmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{pmatrix}$ ,  $\boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{pmatrix}$ ;
  - $\boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{pmatrix}$ .
  - Then  $\mathbf{Y}_1 \sim N_r(\boldsymbol{\mu}_1, \boldsymbol{\Sigma}_{11})$  and  $\mathbf{Y}_2 \sim N_{p-r}(\boldsymbol{\mu}_2, \boldsymbol{\Sigma}_{22})$ .

## Useful properties ii

- Assume the same partition as above. Then the following are equivalent:
  - $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent;
  - $\Sigma_{12} = 0$ ;
  - $\text{Cov}(\mathbf{Y}_1, \mathbf{Y}_2) = 0$ .



## Exercise (J&W 4.3)

Let  $(Y_1, Y_2, Y_3) \sim N_3(\mu, \Sigma)$  with  $\mu = (3, 1, 4)$  and

$$\Sigma = \begin{pmatrix} 1 & -2 & 0 \\ -2 & 5 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

Which of the following random variables are independent?  
Explain.

1.  $Y_1$  and  $Y_2$ .
2.  $Y_2$  and  $Y_3$ .
3.  $(Y_1, Y_2)$  and  $Y_3$ .
4.  $0.5(Y_1 + Y_2)$  and  $Y_3$ .
5.  $Y_2$  and  $Y_2 - \frac{5}{2}Y_1 - Y_3$ .

# Conditional Normal Distributions i

- **Theorem:** Let  $\mathbf{Y} \sim N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ , where
  - $\mathbf{Y} = \begin{pmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{pmatrix}$ ,  $\boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{pmatrix}$ ;
  - $\boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{pmatrix}$ .
- Then the *conditional distribution* of  $\mathbf{Y}_1$  given  $\mathbf{Y}_2 = y_2$  is multivariate normal  $N_r(\boldsymbol{\mu}_{1|2}, \boldsymbol{\Sigma}_{1|2})$ , where
  - $\boldsymbol{\mu}_{1|2} = \boldsymbol{\mu}_1 + \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}(y_2 - \boldsymbol{\mu}_2)$
  - $\boldsymbol{\Sigma}_{1|2} = \boldsymbol{\Sigma}_{11} + \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}\boldsymbol{\Sigma}_{21}$ .

## Conditional Normal Distributions ii

- **Corrolary:** Let  $\mathbf{Y}_2 \sim N_{p-r}(\mu_2, \Sigma_{22})$  and assume that  $\mathbf{Y}_1$  given  $\mathbf{Y}_2 = y_2$  is multivariate normal  $N_r(Ay_2 + b, \Omega)$ , where  $\Omega$  does not depend on  $y_2$ . Then

$$\mathbf{Y} = \begin{pmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{pmatrix} \sim N_p(\mu, \Sigma), \text{ where}$$

- $\mu = \begin{pmatrix} A\mu_2 + b \\ \mu_2 \end{pmatrix};$
- $\Sigma = \begin{pmatrix} \Omega + A\Sigma_{22}A^T & A\Sigma_{22} \\ \Sigma_{22}A^T & \Sigma_{22} \end{pmatrix}.$

## Exercise

- Let  $\mathbf{Y}_2 \sim N_1(0, 1)$  and assume

$$\mathbf{Y}_1 \mid \mathbf{Y}_2 = y_2 \sim N_2 \left( \begin{pmatrix} y_2 + 1 \\ 2y_2 \end{pmatrix}, I_2 \right).$$

Find the joint distribution of  $(\mathbf{Y}_1, \mathbf{Y}_2)$ .

## Another important result i

- Let  $\mathbf{Y} \sim N_p(\mu, \Sigma)$ , and let  $\Sigma = LL^T$  be the Cholesky decomposition of  $\Sigma$ .
- We know that  $\mathbf{Z} = L^{-1}(\mathbf{Y} - \mu)$  is normally distributed, with mean 0 and covariance matrix

$$\text{Cov}(\mathbf{Z}) = L^{-1}\Sigma(L^{-1})^T = I_p.$$

- Therefore  $(\mathbf{Y} - \mu)^T \Sigma^{-1}(\mathbf{Y} - \mu)$  is the sum of *squared* standard normal random variables.
  - In other words,  $(\mathbf{Y} - \mu)^T \Sigma^{-1}(\mathbf{Y} - \mu) \sim \chi^2(p)$ .
  - This can be seen as a generalization of the univariate result  $\left(\frac{X-\mu}{\sigma}\right)^2 \sim \chi^2(1)$ .

## Another important result ii

- From this, we get a result about the probability that a multivariate normal falls within an *ellipse*:

$$P\left((\mathbf{Y} - \mu)^T \Sigma^{-1} (\mathbf{Y} - \mu) \leq \chi^2(\alpha; p)\right) = 1 - \alpha.$$

- We can use this to construct a confidence region around the sample mean.